

## 17. ORTHOPTERA.

### 2. Blattodea

by

R. SHELFORD.

With 2 plates.

---

The number of species of *Blattodea* previously recorded from the region traversed by Dr. YNGVE SJÖSTEDT's expedition is quite insignificant. GERSTÆCKER described in 1869 the collection of insects made by Baron C. C. VON DER DECKEN in the Kilimandjaro district, but scarcely more than a dozen of these were cockroaches. A few species from Mombasa and adjacent localities on the East African coast have been described from time to time by various authors, but in the absence of really representative collections no general review of the East African Blattid fauna has been possible, as in the case of South Africa, Abyssinia, Madagascar, the Cameroons and Angola. This group of insects is usually much neglected by collectors, but this reproach cannot be laid to Dr. SJÖSTEDT's charge, for his collection is a very considerable one, including fifty-one species — about 677 specimens — many of which are represented by long series of individuals of both sexes; in fact the collection may be safely regarded as thoroughly representative of the region in which it was made. The species are referable to 29 genera, four of which are new to science, and of the 51 species, 26 or more than half, are new. In the following account of the collection the species are enumerated and described, species recorded by other authors from the same region — which may be conveniently defined as lying between the equator and 5°S. and between the 35<sup>th</sup> and 40<sup>th</sup> parallels of longitude — are also noted and finally an analysis of this Blattid fauna and its comparison with the corresponding faunas of other regions of Africa is attempted.

Fam. **Ectobiidae.**Gen. **Theganopteryx** BR.**Theganopteryx africana** SAUSS.

*Ectobia africana* SAUSSURE, Abh. Senckenb. Ges. vol. 21 p. 569 (1899)

*Kilimandjaro*: Kibonoto 1000—1300 metres (April); Lower *Meru*: Masai steppes; *Usambara*: Mombo (June); 9 ♂♂, 4 ♀♀.

Previously recorded from the East African coast (VOELTZKOW); the species superficially resembles very closely *Ectobius perspicillaris* HERBST, but structurally differs in the separation of the radial and ulnar veins of the tegmina; since this is the very character used to divide the genus *Theganopteryx* from the genus *Ectobius* the placing of the species in the latter genus is quite illogical.

**Theganopteryx saussurei** SHELF.

*Theganopteryx saussurei* Shelford, Gen. Insect. Fasc. 55. p. 8 (1907).

*Theganopteryx senegalensis* var. Saussure, Ann. Mus. Civ. Gen. Vol. 35 p. 3 (1895).

*Kilimandjaro*: Kibonoto 1300—1900 metres; Lower *Meru*: Masai steppes. 8 ♂♂.

The species is undoubtedly very closely allied to the West African *senegalensis* SAUSS. and may possibly be shown later to be identical; at present however I consider it advisable to separate them.

The two East African species may be separated by their colouration.

- |   |                         |
|---|-------------------------|
| 1. Testaceous, with scattered fuscous points        | <i>africana</i> SAUSS.  |
| 1'. Fuscous, pronotum & tegmina margined with white | <i>saussurei</i> SHELF. |

Gen. **Mallotoblatta** SAUSSURE & ZEHNTNER.**Mallotoblatta kraussi** ADEL.

(Plate 3, fig. 3.)

*Mallotoblatta Kraussi* ADELUNG, Ann. Mus. Acad. Imp. des Sci. St. Petersburg vol. IX. p. 12 (1905).

♂. Flavo-testaceous. Body with a sparse erect pubescence most marked on the ventral surface. A transverse band on the frons, the apex of the maxillary palpi and the labrum fuscous. Vertex of head not covered by pronotum. Pronotum trapezoidal, with two longitudinal fuscous vittae. Tegmina equal to the body in length, 15 costal veins, radial vein bifurcate, 7 longitudinal discoidal sectors, posterior ulnar simple. Wings shorter than the tegmina, mediastinal vein 5-ramose, 9—10 costals, these and the mediastinal rami incrassated, radial vein bifurcate near apex, median vein sinuate, ulnar vein bifurcate near apex, medio-discal and medio-ulnar areas

crossed by transverse nervules, a small triangular apical field. 7<sup>th</sup> abdominal tergite depressed, its posterior border emarginate, the scent-glands opening at the bottom of the depression, 8<sup>th</sup> tergite widely emarginate posteriorly, 9<sup>th</sup> tergite exposed. Supra-anal lamina large, quadrate, posterior angles acute. Cerci elongate. Sub-genital lamina asymmetrical, produced. Front femora with several spines on the anterior margin beneath, the distal ones minute, formula of apical spines  $\frac{2}{1}$ ,  $\frac{1}{1}$ ,  $\frac{1}{1}$ , no genicular spines on the front femora.

Length of body 12 mm; length of tegmina 9,5 mm, pronotum 3 mm  $\times$  3,9 mm.

*Meru*: Rain forest 3,000 m., 1 ♀ (January).

I believe this to be the male of ADELUNG's species described from Abyssinia from a female only. Dr. ADELUNG was not quite certain if this species and *brachyptera* ADEL. also described from a female, were rightly referable to the genus *Mallotoblatta*. The male of *kraussi* certainly does not conform quite strictly to the generic type as represented by *M. pubescens* SAUSS. and ZEHNT., and *M. pilosella* SAUSS. and ZEHNT. from Madagascar, for the pronotum is not »chiffonée», the supra-anal plate is produced and quadrate instead of transverse, the form of the dorsal abdominal tergites and scent-gland opening is different and the front-femora are armed according to the type *A* of SAUSSURE insted of according to type *B*. Nevertheles in the erect pubescence, and in the venation of tegmina and wings *M. kraussi* shows such close affinities to the Mascarene species that for the present it may be allowed a resting-place in the same genus. The discovery of the females of the Mascarene species may occasion the erection of a new genus for the continental forms in which sexual dimorphism is a leading characteristic.

### Gen. *Hololampra* SAUSS.

Two species of this characteristically Palaearctic genus occur in the Kilimandjaro region, one species only is represented by both sexes and the male of this differs from those of all the other representatives of the genus by the length of the wings. The females of the two species may be distinguished as follows:—

1. Testaceous, tegmina and pronotum with brown points . . . . . *sjöstedti* n. sp.
- 1'. Abdomen and head piceous . . . . . *aethiopica* n. sp.

### *Hololampra aethiopica* n. sp.

(Plate 2, Fig. 1.)

♂. Piceous. Antennae fuscous. Pronotum trapezoidal, margins broadly hyaline, disc with scattered brown points or with dark castaneous blotches symmetrically arranged. Scutellum exposed. Tegmina equal to the body in length, clear testaceous with scatted castaneous points on the veins and minute brown maculations between the veins; 9–10 costal veins, 5–6 oblique discoidal sectors, anal vein impressed. Wings abbreviated, reaching the 6<sup>th</sup> abdominal tergite, hyaline veins infuscated,

5—6 rudimentary costal veins, distal half of median vein alone visible, medio-discal area crossed by transverse nervules, ulnar vein simple, 1<sup>st</sup> axillary triramous. Abdomen above with the first 6 abdominal tergites piceous, the remainder flavo-testaceous, the posterior borders of 6<sup>th</sup>—8<sup>th</sup> tergites emarginate, opening of scent-glands on 7<sup>th</sup> tergite, supra-anal lamina transverse, slightly produced—Cerci fuscous, 7 jointed. Abdomen beneath piceous, laterally margined with testaceous; sub-genital lamina irregular, elongate.

Femora castaneous, sparsely armed beneath, no genicular spines on the middle pair; tibiae and basal joints of tarsi flavo-testaceous, apical joints of tarsi fuscous.

♀. Larger; variable in colour, pronotum with brown points or heavily blotched with castaneous. Tegmina extending to 1<sup>st</sup> or 2<sup>nd</sup> abdominal tergite. Wings rudimentary. Abdomen piceous above, with the tergites margined with flavo-testaceous, or with flavo-testaceous spots only on their lateral margins; supra-anal lamina short, transverse; abdomen beneath piceous, laterally margined with testaceous, sub-genital lamina ample. Cerci piceous. Legs as in ♂.

Larvae have all the thoracic tergites testaceous variegated with castaneous, the abdomen piceous; the legs are entirely castaneous.

♂. Total length 6 mm, length of tegmina 5,9 mm; pronotum 2 mm × 3 mm.  
 ♀. » » 7,1 » » » 3 » » 2,2 » × 3,2 »

*Kilimandjaro*: Rain forest, Kiboseho, 3,000 m (February), 2 ♂♂, 6 ♀♀, 5 larvae.

#### *Hololampra sjöstedti* n. sp.

♀. Testaceous. Head rufo-testaceous, vertex covered. Pronotum trapezoidal, sides deflexed, lateral margins broadly hyaline, posterior border truncate, exposing the scutellum. Tegmina lanceolate, attaining the sixth abdominal segment, overlapping slightly, 8 costal veins, one ulnar vein with 5 branches, anal vein impressed, reaching the sutural margin at one half of its length; a few brown points occur on the veins. Wings rudimentary. Abdomen above variegated with fuscous, supra-anal lamina short transverse; abdomen beneath variegated with fuscous, disc fuscous, sub-genital lamina ample, its centre rufo-testaceous or castaneous; cerci six-jointed, rufo-testaceous. All the femora with two spines only on the anterior margin beneath, none on the posterior margin, with genicular spines and two apical spines.

Total length 6,5 mm; length of tegmina 4 mm; pronotum 2 mm × 3 mm.

*Kilimandjaro*: Kibonoto, zone of culture 1,000—1,900 m (August to December).

Lower *Meru*: Ngare na nyuki (January); Masai steppes (October). 22 ♀♀, 2 larvae. One example has the ootheca projecting from the end of the abdomen; it is chitinous and carried with the suture uppermost, it is not earinate.

The species appears to be nearest to *H. minuta* SHELF. from Madagascar.

Fam. **Phyllodromiidae.**Gen. **Ischnoptera** BURM.

The East African species can readily be distinguished from species of *Phyllodromia*, not only by the wing venation but also by the longitudinal, instead of oblique, discoidal sectors of the tegmina and by the asymmetry of the sub-genital lamina and styles in the male sex. The following table will assist in the determination of the known species:—

1. Testaceous or rufo-testaceous species.
  2. Scent-glands of ♂ opening at base of supra-anal lamina, which is not much produced . . . . . *bimaculata* GERST.
  - 2'. Scent-glands of ♂, not as above, supra-anal lamina produced *strigosa* SCHAUM.
- 1'. Rufo-castaneous or rufo-fulvous species.
  2. Small . . . . . *neutra* SAUSS.
  - 2'. Larger . . . . . *incuriosa* SAUSS.

*I. strigosa* SCHAUM. is recorded from Mosambique, it cannot be recognized with certainty. *I. neutra* SAUSS. from »Africa meridionalis» is described from a female only and appears to differ in size only from *I. incuriosa* SAUSS.

**Ischnoptera bimaculata** GERST.

(Pl. 3, figs. 10 &amp; 15.)

*Phyllodromia bimaculata* GERSTAECKER, Arch. Naturg. XXXV, p. 206 (1869);  
 Von der Decken, Reisen in Ost-Afrika III (2) p. 4 (1873).

GERSTAECKER'S description does not include an account of the wing structure in this species; the venation is somewhat variable but at least two branches of the ulnar vein fail to reach the outer margin of the wing and are directed towards the dividing vein. On account of this character the species must be placed in the genus *Ischnoptera*. The 8<sup>th</sup> abdominal tergite in the male is completely hidden beneath the 7<sup>th</sup> tergite, the 9<sup>th</sup> tergite is also invisible and the base of the supra-anal lamina is depressed forming a cavity fringed with hairs, the scent glands open on the sides of two chitinous tubercles in this cavity. The sub-genital lamina is very irregular and notched on the left side, a style being situated in the notch, the right style is stouter and spinose, the apex of the lamina forms a hirsute lobe. Ootheca as in *Phyllodromia germanica* L.

*Kilimandjaro*: Kibonoto, lower slopes and 1,300—1,900 mètres, under dead leaves in banana plantations. 21 ♂♂, 25 ♀♀. The species was originally recorded from Lake Jipe at Kilimandjaro (VON DER DECKEN).

Var. **sobrina** nov.

♂ and ♀. Castaneous, margins of pronotum hyaline its disc castaneous. Wings with veins fuscous.

1 ♂, 21 ♀♀ and numerous larvae. From the same localities as the typical form and also from Usambara. Structurally these two forms are identical and of similar size.

**Ischnoptera incuriosa SAUSS.**

(Pl. 3, figs. 6 & 7.)

*Ischnoptera incuriosa* DE SAUSSURE, Abh. Senckenb. Ges. XXI. p. 571 (1899).

*Kilimandjaro*: Kibonoto 1,000—1,300 mètres (November and December), 2 ♂♂. Originally described from East-Africa e coll. VOELTZKOW.

Gen. **Phyllodromia SERV.**

Key to the East-African species.

1. ulnar vein of wings simple or bifurcate . . . . . *germanica* L.
- 1'. ulnar vein of wings multiramose.
  2. pronotum with two longitudinal fuscous vittae . . . *bivittata* SERV.
  - 2'. pronotum not as above.
    3. tegmina with numerous castaneous points . . . . *zehntneri* nom. n.
    - 3'. tegmina not as above.
      4. Opening of ♂ scent-glands situated on 7<sup>th</sup> abdominal tergite.
        5. Sub-genital lamina in ♂ with margins inflected, styles flattened . . . . . *supellectilium* SERV.
        - 5'. Sub-genital lamina in ♂ not as above, styles not flattened.
          6. Frons flattened . . . . . *nigromarginata* sp. n.
          - 6'. Frons rounded . . . . . *sjöstedti* n. sp.
      - 4'. Opening of ♂ scent-glands not situated on 7<sup>th</sup> abdominal tergite.
        5. Opening of ♂ scent-glands not visible . . . *insignis* sp. n.
        - 5'. Opening of ♂ scent-glands situated on supra-anal lamina. . . . . *testacea* sp. n.

*P. trigonalis* SAUSSURE from »Africa meridionalis» (VOELTZKOW) was described from a female; the rufous colour, trigonal supra-anal lamina, and front femora with completely armed anterior margin beneath, appear to be the distinctive features of the species.

**Phyllodromia germanica L.**

*Blatta germanica*, LINNÆUS Syst. Nat. (ed. 12) I (2) p. 668 n. 7 (1767).

*Kilimandjaro*: Kibonoto, 1,000—1,300 mètres; *Meru* rain-forest; Massai steppes (January & May). 1 ♂, 7 ♀♀, 2 larvae. A cosmopolitan species.

*Phyllodromia bivittata* SERV. is not represented in Dr SJÖSTEDT's collection but has been recorded from Wanga (VON DER DECKEN); it is a cosmopolitan species and

can be distinguished from *P. germanica* L., by the castaneous stripe on the tegmina, the ramose ulnar vein of the wings, the notched supra-anal lamina of the male and transverse supra-anal lamina of the female and by the produced, and laterally compressed sub-genital lamina of the male, with finely cleft apex.

***Phyllodromia supellectilium* SERV.**

(Pl. 3, fig. 11.)

*Blatta supellectilium* SERVILLE, Hist. Ins. Orth. p. 114 (1839).

*Blatta extenuata* WALKER, Cat. Blatt. Brit. Mus. p. 221 (1868).

*Blatta figurata* WALKER, Cat. Derm. Salt. Brit. Mus. V. Suppl. Blatt. p. 24 (1871).

*Blatta transversalis* WALKER, Cat. Derm. Salt. Brit. Mus. V. Suppl. Blatt. p. 25 (1871).

*Phyllodromia delta* KIRBY, Ann. Mag. Nat. Hist. (7) V. p. 280 (1900).

*Kilimandjaro*: Kibonoto 1,000—1,300 mètres; Massai steppes. Lower *Meru*.  
*Usambara*: Mombo. 5 ♂♂. A cosmopolitan species, which is very variable in size; it can however always be recognized by the form of the sub-genital plate of the male. The lateral margins of this are inflected and the two styles are flattened. The supra-anal lamina is almost hidden by the ninth tergite; in the centre of the seventh tergite is a circular depression from which rises a bifurcate chitinous structure covered with a fine pubescence and the scent-glands open on either side of this.

Dr. SJÖSTEDT's specimens have been compared with three examples in the Oxford Museum determined by SERVILLE himself and though the Kilimandjaro specimens are larger and darker than SERVILLE's specimens (= *P. delta* KIRBY), I have no hesitation in referring them to *supellectilium* on account of the identity of structure. The female of *P. supellectilium* is very different from the male and this sexual dimorphism is a character that perhaps entitles this species to generic rank separate from *Phyllodromia*.

***Phyllodromia zehntneri* nom. nov.**

*Theganopteryx (Pseudectobia) punctulata* DE SAUSSURE & ZEHNTNER, Grandidier. Hist. Madagasc. Orth. I p. 15 (1895).

*Usambara*: Tanga (June). 1 ♀.

I now think that I was wrong in referring this species to the genus *Theganopteryx*, sens. strict. (cf. Ann. Mag. Nat. Hist. (7) XIX. p. 36. 1907); the ulnar vein of the wing is described as »bi-ramose» not as »bifurcate» and its branches vary in number from three to four. The species also can not be referred to the genus *Pseudectobia* as defined by me (l. c.) but appears to fall quite naturally into the genus *Phyllodromia*. Unfortunately it is necessary to find a new specific name for it, since *punctulata* has already been applied to two species of *Phyllodromia*, viz. *P. punctulata* BEAUVOIS (1805) and *P. punctulata* BRUNNER (1893), for the latter species the name *brunneri* is suggested. *P. zehntneri* was previously recorded from Madagascar.

***Phyllodromia nigromarginata* sp. n.**

(Pl. 3, fig. 12.)

♂. Testaceous. Head castaneous, vertex not covered by pronotum flattened, antennae testaceous. Pronotum transversely elliptical, variegated with castaneous,

with two oblique sulci on disc, lateral margins hyaline. Tegmina testaceous, 10 to 11 costals, the last 3 ramose, anterior ulnar sending 5 oblique branches to the sutural margin and 2 ramose branches to the apical margin, posterior ulnar simple. Wings hyaline, mediastinal vein unbranched, 10 costals, their apices incrassated, ulnar vein with 5 branches. Abdomen above with the disc testaceous with a central piceous blotch, margins castaneous, the last 4 tergites piceous; the scent glands open on the 7<sup>th</sup> tergite by two orifices, a papilla covered with fine setae placed between the openings; 9<sup>th</sup> tergite concealed; supra-anal lamina trigonal. Abdomen beneath rufo-testaceous broadly margined with dark castaneous; sub-genital lamina produced, no styles; cerci fuscous. Legs testaceous, front femora armed on the anterior margin beneath with a complete row of spines, the proximal spines being the longer; each femur with 1 genicular and 2 apical spines.

Total length 16 mm.; length of body 11,2 mm.; length of tegmina 14 mm.; pronotum 2,9 mm. × 4,2 mm.

Lower *Meru*: Ngare na nyuki. *Kilimandjaro*: Kibonoto 1,000—1,900 mètres, 6 ♂♂.

*Phyllodromia sjöstedti* sp. n.

♂. Closely allied to the preceding species but smaller. Rufo-testaceous. Head castaneous, vertex rounded, not covered by pronotum; antennae infuscated, testaceous at base. Pronotum transversely elliptical, disc castaneous with a rufo-testaceous, macula on the posterior part, lateral margins broadly hyaline. Tegmina rufo-castaneous 11 costals, the last two ramose, anterior ulnar sending ramose branches to sutural and apical margins. Wings hyaline, suffused with rufo-testaceous on anterior margin, mediastinal vein simple, 9 costals the first 6 incrassated, ulnar vein with 5 branches. Abdomen above castaneous, scent-glands as in preceding species, supra-anal lamina shortly trigonal; abdomen beneath castaneous with the disc rufo-testaceous, sub-genital lamina ample, produced, slightly pubescent, with two styles; cerci elongate, fuscous. Legs testaceous, front femora armed beneath on anterior margin with a complete row of spines, of which the more distal are the longer; formula of apical spines  $\frac{1}{1}$ ,  $\frac{1}{1}$ ,  $\frac{1}{1}$ .

Total length 14 mm.; length of body 10,6 mm.; length of tegmina 11,8 mm.; pronotum 3 mm. × 4,2 mm.

Lower *Meru* (November), 1 ♂.

*Phyllodromia insignis* sp. n.

(Pl. 3, fig. 8.)

♂. Castaneous. Antennae fuscous. Vertex of head not covered by pronotum. Pronotum transversely elliptical, lateral margins broadly hyaline, disc with some obscure impressions. Tegmina with costal margin to near apex hyaline, remainder of tegmina not uniformly castaneous but internervular spaces crossed by numerous minute hyaline streaks arranged more or less regularly; 9 costal veins, radial bifurcate and sending branches both to costal margin and to apex, anterior ulnar with 6 oblique branches, posterior ulnar simple. Wings infuscated, mediastinal vein with 2 branches, 8 costals, their apices incrassated, ulnar vein tri-ramose, the branches bifurcate.



Abdomen with the disc above testaceous; supra-anal lamina transverse; sub-genital lamina cucullate, deeply notched, the flattened styles springing from the borders of the notch; cerci elongate, fuscous. Legs testaceous, front femora armed with piliform setae only on the anterior margin beneath.

Total length 12 mm.; length of tegmina 9,5 mm.; pronotum 3 mm.  $\times$  3,9 mm.

*Kilimandjaro*: Kibonoto 1,000—1,300 mètres (Sept.), 1 ♂.

#### *Phyllodromia testacea* sp. n.

(Pl. 2, fig. 14.)

♂. Pale testaceous. Head rufo-testaceous, vertex not covered by pronotum. Pronotum transversely elliptical, lateral margins broadly hyaline. Tegmina with marginal area very broad, 9 costal veins, the last ramose, anterior ulnar sending 5 branches to sutural and apical margins, the latter ramose; numerous transverse venulae between the veins. Wings hyaline, mediastinal vein bifurcate, the lower branch bi-ramose, 6 costal veins their apices not incrassated, end of radial vein ramose, ulnar vein with 4 branches. Abdomen rufo-testaceous towards apex, supra-anal lamina transverse, posteriorly incrassated, a depression occurs in the posterior edge on either side of the middle line and at the base of these the scent glands open. Sub-genital lamina produced considerably beyond the supra-anal lamina, cucullate at apex which is deeply and narrowly cleft, the flattened styles spring from the posterior margin; cerci long, testaceous. Front femora with piliform setae only on the anterior margin beneath; all the femora with genicular spines; formula of apical spines  $\frac{2}{1}$ ,  $\frac{1}{1}$ ,  $\frac{1}{1}$ .

Total length 14 mm.; length of body 11 mm.; length of tegmina 12 mm.; pronotum 3,6 mm.  $\times$  5 mm.

*Usambara*: Tanga (June), 1 ♂.

The nearest allies of this species appear to be *P. laterifera* Wlk., *P. propinqua* Wlk., *P. majuscula* Wlk. from the Indo-Malayan and Indo-Australian regions of the world. The remarkable form of the supra-anal lamina of *P. testacea* is however a sufficiently distinctive character and should render it easier of identification than is the case with so many of the obscure species of this large genus.

#### Gen. *Ceratinoptera* BRUNNER.

This genus should be reserved for those species of Phyllodromiinae characterized by short or reduced tegmina and short or rudimentary wings, the tegmina when reduced are lanceolate, not truncate nor lobiform, and the wings when present have the ulnar vein bifurcate or simple. In the genus *Allacta* the wings are well-developed but the ulnar vein is ramose, the median is sometimes absent. The genus *Temnopteryx* is characterized — as its name signifies — by the truncate tegmina and rudimentary wings.

The African species of *Ceratinoptera* may be distinguished as follows:

1. Castaneous species.
2. Tegmina not banded nor variegated with paler colour.
  3. Pronotum not pale-bordered anteriorly.
  4. Tegmina much shorter than the body.
    5. Sub-genital lamina of ♂ posteriorly truncate, angles acute . *abbreviata* SAUSS. (Réunion).
    - 5'. Sub-genital lamina of ♂ posteriorly rounded . . . . . *castanea* sp. n. (E. Africa).
  - 4'. Tegmina not or scarcely shorter than the body.
    5. Larger, legs testaceous . . . . *madecassa* SAUSS. (Madagascar).
    - 5'. Smaller, legs castaneous . . . . *orata* sp. n. (E. Africa).
  - 3'. Pronotum pale-bordered anteriorly . *abyssinica* SAUSS. (Abyssinia).
- 2'. Tegmina banded or variegated with paler colour.
  3. Pronotum with disc rufous.
    4. Minute species . . . . . *perpulchra* sp. n. (E. Africa).
    - 4'. Larger . . . . . *dimidiata* GERST. (E. Africa).
  - 3'. Pronotum with disc not rufous.
    4. Disc of pronotum with testaceous maculae . . . . . *variegata* SCHULTH (E. Africa)  
 = *hottentota* SAUSS. (Delagoa Bay)  
 = *transvaaliensis* KIRBY (Transvaal).
    - 4'. Disc of pronotum concolorous . . *bolivari* ADEL. (Gallaland).
- 1'. Testaceous or ferruginous species.
  2. Pronotum with fuscous markings on disc.
    3. Markings on disc of pronotum numerous . . . . . *inscripta* WLK. (Natal).
    - 3'. Markings on disc reduced to two obsolescent vittae . . . . . *bimaculata* sp. n. (E. Africa).
  - 2'. Pronotum without markings on disc.
    3. Supra-anal lamina of ♂ transverse . *ferruginea* SCHULTH (S., W. & E. Africa).
    - 3'. Supra-anal lamina of ♂ produced.
      4. Opening of scent-glands at base of supra-anal lamina . . . . . *variabilis* sp. n. (E. Africa).
      - 4'. Opening of scent-glands on 7<sup>th</sup> abdominal tergite . . . . . *sjöstedti* sp. n. (E. Africa).

*Blatta misella* STÅL. from Natal is a species of *Hololampra*.

*Ceratinoptera bimaculata* sp. n.

♀. Testaceous, broad, short. Frons with a castaneous band; apical joint of maxillary palpi and antennae except at base, fuscous. Vertex of head almost covered by the pronotum. Pronotum large, trapezoidal, lateral margins broadly, posterior margin narrowly hyaline, posterior border slightly obtusely angled, scutellum not exposed; two obsolescent castaneous vittae on the disc. Tegmina longer than the body, testaceo-hyaline, membranous, 10 costal veins, ulnar, anal and axillary veins obsolescent. Wings abbreviated, extending to the 2<sup>nd</sup> abdominal tergite. Abdomen above castaneous, laterally margined with testaceous; supra-anal lamina testaceous, small, triangular. Abdomen beneath testaceous, with lateral fuscous maculae obsolescent posteriorly; sub-genital lamina very large, semiorbicular. Legs testaceous, front femora with a complete row of long spines on the anterior margin beneath.

Total length 9.1 mm.; length of body 8 mm.; length of tegmina 6.1 mm.; pronotum 3 mm. × 4 mm.

*Usambara*: Mombo (June), 2 ♀♀.

The broad abdomen and the obsolescent venation of the membranous tegmina are the characteristic features of this species. Obsolescent venation is usually associated with a corneous and more or less opaque texture of the tegmina, its disappearance from delicate membranous tegmina is most unusual.

*Ceratinoptera castanea* sp. n.

♂. Allied to *C. abbreviata* SAUSS. and *C. madecassa* SAUSS. but differing from the former by its larger size and by the different shape of the sub-genital lamina and from the latter by the shorter tegmina.

Dark castaneous. Antennae fuscous, mouth parts testaceous. Pronotum trapezoidal, nearly covering vertex of head lateral margins rufo-castaneous, its posterior margin slightly obtusely angled. Tegmina lanceolate, corneous, extending to the 4<sup>th</sup> abdominal tergite, 10 costals, anal vein impressed, remainder of venation obscured. Wings much reduced, extending to middle of 2<sup>nd</sup> abdominal tergite. Supra-anal lamina transverse, slightly produced. Sub-genital lamina broad, transverse, slightly produced, its posterior angles rounded, with two short styles. Legs rufo-testaceous.

Length of body 12 mm.; length of tegmina 7 mm.; pronotum 4 mm. × 5 mm.

*Usambara*: Mombo (June), 1 ♂.

*Ceratinoptera sjöstedti* sp. n.

(Pl. 3, figs. 16—17.)

♂. Allied to *C. ferruginea* SCHULTZ but with longer tegmina. Rufo-testaceous. Head castaneous, antennae testaceous at base, remainder infuscated; vertex of head not nearly covered by pronotum. Pronotum transversely elliptical with lateral margins hyaline, posterior border truncate, exposing the scutellum. Tegmina extending to penultimate segment, membranous; marginal field very broad, 12 costal veins, anterior

ulnar with 6 branches, posterior ulnar simple, 5 axillary veins. Wings reduced, attaining 2<sup>nd</sup> abdominal tergite. Opening of scent-glands a circular orifice on the 7<sup>th</sup> abdominal tergite. Supra-anal lamina trigonal, posterior border slightly emarginate: sub-genital lamina broad, produced, apex slightly cleft, with two styles asymmetrically placed. Titillator spinous.

Length of body 8,2 mm.; length of tegmina 7,5 mm.; pronotum 3 mm. × 4 mm.  
Lower *Meru*: at the river Ngare na nyuki (January), 2 ♂♂.

*Ceratinoptera variabilis* sp. n.

♂. Rufo-testaceous. Head with a castaneous band between the eyes and two spots between the antennal sockets; antennae testaceous at base; vertex of head not covered by pronotum. Pronotum trapezoidal, sides deflexed, posterior margin obtusely angled, margins hyaline, disc rufo-testaceous or castaneous variegated with rufo-testaceous. Tegmina lanceolate, reaching the 7<sup>th</sup> segment, mediastinal area hyaline, mediastinal vein bi-ramose, 6 costals the last ramose, discoidal field with 5 longitudinal sectors. Wings minute, extending to 1<sup>st</sup> abdominal tergite. Abdomen rufo-testaceous variegated with castaneous, 7<sup>th</sup> tergite cucullate & concealing the 8<sup>th</sup> & 9<sup>th</sup> tergites; the scent-glands open at the base of the supra-anal lamina which is coveate at its base with a median carina, the opening of the glands is fringed with hairs; supra-anal lamina produced, sub-quadrate, not exceeding the sub-genital lamina which is asymmetrical, terminating at apex in a rounded hirsute lobe and with two unequal styles. Front femora on anterior margin beneath armed with a complete row of spines, the more proximal the longer.

♀. Castaneous. Lateral margins of pronotum hyaline, disc sometimes with two or more testaceous maculae, scutellum exposed. Tegmina abbreviated, obovate, not extending beyond the 4<sup>th</sup> abdominal tergite, 9 costals, anterior ulnar with 4 oblique branches, posterior ulnar simple, anal vein impressed reaching sutural margin at a point one-third from the apex. Wings extending to third abdominal tergite, apex infuscated, costal veins obsolete, median vein straight prominent, ulnar vein stout, bifurcated at apex, 1<sup>st</sup> axillary vein 3-ramose, stout, the first branch with 3 short branches at apex. Abdomen piecous above and below; supra-anal lamina triangular; apex emarginate, sub-genital plate ample, semi-orbicular.

♂. Length of body 9,2 mm.; length of tegmina 5,2 mm.; pronotum 3,1 mm. × 4 mm.

♀. » » » 11 mm.; » » » 5,5 mm.; » 3 mm. × 4,2 mm.

*Kilimandjaro*: Kibonoto 1,000—3,500 mètres (Sept.—Nov.), 4 ♂♂, 3 ♀♀, 8 larvae.

The apex of the abdomen is very similar in construction to that of *Ischnoptera bimaculata* GERST.

♂. var. *truncata* nov.

Similar to above but tegmina and wings shorter, the former transversely truncate. 1 ♀. Lower *Meru*.

This low country specimen may, possibly be a distinct species, but in colour and general appearance it resembles so closely the mountain forms that I hesitate to separate it, at any rate until a male from the same habitat is found.

*Ceratinoptera perpulehra* sp. n.

(Pl. 2, fig. 2; Pl. 3, fig. 13.)

♂. Minute, rufo-testaceous, nitid. Head castaneous, rufous on vertex, basal joints of antennae testaceous, remainder fuscous; vertex covered by pronotum. Pronotum trapezoidal with two longitudinal vittae castaneous. Tegmina coriaceous, just failing to reach end of abdomen, a broad humeral stripe which extends as two narrow lines along the radial and ulnar veins, castaneous; 12 costal veins, anterior ulnar bi-ramose, the remaining veins not visible. Wings rudimentary. Abdomen above rufo-testaceous with lateral margins and a central patch castaneous; scent-glands opening on 7<sup>th</sup> tergite, at base of two deep depressions; supra-anal lamina produced, rounded; cerci short, rufo-testaceous; abdomen beneath castaneous, sub-genital lamina rufo-testaceous, rounded, produced, styles asymmetrically placed. Legs testaceous, front femora as in preceding species.

♀. Somewhat larger, dark castaneous, the disc of the pronotum and field of tegmina rufous, lateral margins of pronotum and mediastinal field of tegmina pale testaceous; abdomen entirely castaneous, cerci fuscous. Tegmina shorter, wings rudimentary, supra-anal lamina triangular, sub-genital lamina ample, semi-orbicular. Legs testaceous, except the bases of the coxae which are castaneous.

♂. Length of body 5,5 mm.; length of tegmina 4 mm.

♀. » » » 7 mm.; » » » 3,8 mm.

*Kilimandjaro*: Kibonoto 1,000—1,900 mètres (March to May, Sept., Oct.), 5 ♂♂, 5 ♀♀, 1 larva.

This is quite the smallest species of the genus; in colouration it approaches *C. dimidiata* GERST. but it lacks the transverse band on the tegmina.

*Ceratinoptera ovata* sp. n.

♀. Convex, castaneous, nitid. Vertex of head not covered by pronotum, antennae piceous. Pronotum trapezoidal, sides deflexed, posterior margin obtusely angled, lateral margins paler than the disc. Tegmina reaching base of supra-anal lamina, mediastinal area testaceous, 14 costals, anal vein impressed, remaining veins obscure, the part of the right tegmen overlapped by the left reticulated. Wings shorter than tegmina, 8 costals, their apices slightly incrassated, the last and apex of radial vein tri-ramose, ulnar vein bifurcated. Abdomen and cerci piceous, supra-anal lamina triangular, sub-genital lamina ample, semi-orbicular. Legs rufous.

Length of body 9 mm.; length of tegmina 6,4 mm.; pronotum 3 mm. × 3,9 mm.

*Usambara*, 1 ♀.

*C. dimidiata* GERST. is probably the nearest ally of the species but it was described from an imperfect specimen so that certainty on this point is impossible.

*Ceratinoptera dimidiata* GERST. has been recorded from Endara, E. Africa (VON DER DECKEN).

### Gen. **Temnopteryx** BRUNNER.

Key to the East African species.

1. Fuscous, pronotum margined with testaceous . . . . . *abyssinica* SAUSS. & ZEHNT.
- 1'. Rufous or testaceous.
  2. Testaceous.
    3. Larger, 9 mm. long . . . . . *ectobioides* sp. n.
    - 3'. Smaller, 7 mm. long . . . . . *affinis* sp. n.
  - 2'. Rufo-testaceous or rufous.
    3. Pronotum with central testaceous vitta . . . . . *caffra* SAUSS.
    - 3'. Pronotum unicolorous . . . . . *rufa* sp. n.

#### **Temnopteryx abyssinica** SAUSS. & ZEHNT.

*Temnopteryx abyssinica* SAUSSURE & ZEHNTNER, Grandidiers Hist. Madagascar Orth. I p. 51 (1895)<sup>1</sup>.

*Temnopteryx saussurei* BOLIVAR, Ann. Soc. Ent. France, vol. LXVI p. 292 (1897).

*Kilimandjaro*: Kibonoto 1,000—3,500 mètres; Masai steppes.

Lower *Meru* (Sept. to Oct.); 10 ♀♀, 5 larvae.

These specimens differ very slightly in colour from the type, which occurs at Massowa; the posterior border of the pronotum is narrowly testaceous, and not provided with a testaceous macula as in the Abyssinian examples, the cerci are testaceous instead of fuscous, and the supra-anal lamina is fuscous instead of testaceous; with these slight differences excepted, the Kilimandjaro examples appear to be identical with the typical specimens. In one example the egg-mass is protruding from the cloaca, the eggs are enclosed in a thin transparent membrane through which the eyes of the developing embryos can be distinctly seen; it is probably the case that this species carries the eggs until they are almost ready to hatch out. Another example is stated to have been found with termites, but the association was probably accidental.

#### **Temnopteryx ectobioides** sp. n.

(Pl. 2, fig. 12.)

♂. Pale testaceous. Head with a fuscous mark between the eyes and another between the antennal sockets, the area between these two marks, pale pinkish-white and nitid; antennae testaceous at base, remainder fusco-testaceous. Pronotum very broad, transversely truncate behind, just failing to cover vertex of head in front; lateral margins broadly hyaline, disc slightly rugulose with a few brown points and with a short fuscous vitta at each posterior angle. Tegmina quadrate just covering the first abdominal segment, with a few brown points between the veins, anal vein

<sup>1</sup> *Temnopteryx abyssinica* SAUSS. (Mém. Soc. Genève. XXIII p. 93. 1873) having been removed to the genus *Ceratinoptera* (vide antea), this species may be allowed to retain its original name.

not marked. Wings minute. Abdomen variegated with rufous and fuscous, more heavily marked below; cerci testaceous spotted with fuscous; supra-anal lamina triangular, apex notched; sub-genital lamina large, trigonal, a median carina, apex deeply cleft, two symmetrically disposed styles, which are bent downwards at right angles to the plane of the lamina. Legs testaceous; front femora armed on the anterior margin beneath with minute piliform spines; formula of apical spines,  $\frac{1}{1}$ ,  $\frac{1}{1}$ ,  $\frac{1}{1}$ .

♀. Darker than male. Head similar but with a triangular castaneous spot at base of clypeus. Pronotum with an incomplete fuscous vitta on each side of the disc and continued on the meso- and metanotum; metanotum with a central fuscous vitta. Tegmina as in the male. Abdomen almost entirely dark castaneous variegated with testaceous; cerci fuscous, tipped with testaceous; supra-anal lamina triangular, apex emarginate; sub-genital lamina ample, semi-orbicular, posterior margin triangularly notched.

♂. Length of body 9 mm.; length of tegmina 2,1 mm.; pronotum 2,9 mm. × 4,8 mm.

♀. " " " 9 mm.; " " " 3 mm. × 4,6 mm.

3 ♂♂, 2 ♀♀. Lower *Meru* (December).

The nearest allies of the species are *T. nana* SAUSS. from Senegal and *T. brachyptera* BOL. from North Africa, from both it may be distinguished by the form of the supra-anal and sub-genital plates.

#### *Temnopteryx affinis* sp. n.

♀. Allied to the preceding species, but smaller and entirely pale testaceous with only a few brown points on prothorax and tegmina. Tegmina quadrate, not extending beyond metanotum, mediastinal vein nearly reaching outer posterior angle, 4 costals, anal vein not marked. Abdomen above with four obscure longitudinal fuscous vittae, supra-anal lamina slightly produced, rounded, sub-genital lamina semi-orbicular, ample.

Length of body 7,2 mm.; length of tegmina 2 mm.; pronotum 2 mm. × 3,5 mm.

*Meru* rain-forest, 3,000—3,500 mètres; 1 ♀.

#### *Temnopteryx rufa* sp. n.

♀. Rufous, nitid. Vertex of head not covered by pronotum; antennae fuscous, except basal joints which are rufous. Pronotum with sides deflexed, posterior margin very slightly angulated. Tegmina truncate, reaching 2<sup>nd</sup> abdominal segment, mediastinal vein sending two branches to outer margin, 10 costals, marginal field very broad, discoidal field narrow, anal vein nearly reaching apex of tegmina. Wings a little shorter than tegmina. Abdomen rufo-castaneous, supra-anal lamina trigonal produced, cerci castaneous, sub-genital lamina ample. Legs rufo-testaceous, front femora with anterior margin beneath armed throughout with spines, the distal shorter than the proximal; all the femora with genicular spines, formula of apical spines  $\frac{1}{1}$ ,  $\frac{1}{1}$ ,  $\frac{1}{1}$ .

Total length 13 mm.; length of tegmina 5 mm.; pronotum 4 mm. × 5,2 mm.

*Kilimandjaro*: Kibonoto 1,300—1,900 mètres (November); 1 ♀.

*T. phalerata* Sss. from S. Africa is perhaps the nearest ally of this species, which however can be distinguished by its smaller size and different colour.

### Gen. **Loboptera** Br.

#### **Loboptera nitida** sp. n.

♀. Rufo-castaneous. Antennae with basal half testaceous, apical half fuscous. Vertex of head just covered by the pronotum. Pronotum transversely elliptical, posteriorly truncate, lateral margins broadly hyaline; tegmina hyaline, lobiform; lateral margins of metanotum hyaline. Abdomen broad, above castaneous, variegated with testaceous, beneath rufo-testaceous, darker on the sides; penultimate tergite with posterior margin produced in the middle, supra-anal lamina triangular, apex slightly carinate, emarginate, sub-genital lamina ample; cerei rufo-testaceous. Legs testaceous: front femora armed on the anterior margin beneath with piliform spines, formula of apical spines  $\frac{2}{1}$ ,  $\frac{1}{1}$ ,  $\frac{1}{1}$ , no genicular spine on front femora.

Length of body 8 mm.; length of tegmina 1,6 mm.; pronotum 3,8 mm.  $\times$  4 mm.

*Usambara*: Mombo (Jan. and June), 2 ♀♀.

One of the two specimens is very much darker than the other, being entirely castaneous both above and beneath.

The species recorded by Adelung from Abyssinia, *L. ras* has no arolia between the tarsal claws and therefore falls into the genus *Paraloboptera* SAUSS.

### Gen. **Apteroblatta** nov.

Allied to *Loboptera* Br. but tegmina entirely absent in both sexes. Eyes rather small, vertex very broad. Supra-anal lamina in both sexes triangular, produced, sub-genital lamina of male slightly produced with two symmetrical styles. Ootheca chitinous, carried with the suture uppermost. Front femora with anterior margin beneath with piliform setae, hind femora heavily armed with spines. Arolia present between tarsal claws.

#### **Apteroblatta perplexa** sp. n.

(Pl. 2, fig. 3 and 13.)

♂. Rufo-testaceous. Vertex of head not covered by pronotum. Thoracic tergites variegated with rufo-castaneous. Abdomen above with disc dark castaneous, marginal castaneous spots on tergites 2—6. Cerei castaneous. Abdomen beneath rufo-testaceous; sub-genital lamina with apex cleft. Legs testaceous; all the femora with genicular spines; formula of apical spines  $\frac{1}{0}$ ,  $\frac{1}{1}$ ,  $\frac{1}{1}$ .

♀. Similar to ♂ but larger. Castaneous markings on thoracic tergites forming a definite pattern. Abdomen above castaneous with a submarginal rufo-testaceous vitta on either side, not extending beyond 6<sup>th</sup> tergite, terminal tergites castaneous; supra-anal lamina with apex slightly notched. Cerei testaceous. Abdomen beneath rufo-castaneous, sub-genital lamina ample, semi-orbicular. Ootheca as in *Loboptera*.



♂. Length 6 mm.; pronotum 2 mm. × 2,5 mm.

♀. » 7—8 mm.; » 2,2 mm. × 3,1 mm.

*Kilimandjaro*: Kiboscho, » Bergwiesen », 3,000 mètres. *Meru* rain-forest, 3,500 m.  
5 ♂♂ and a large number of ♀♀.

Were it not for the presence in the series, of female specimens with the ootheca protruding from the cloaca, these minute cockroaches might readily be overlooked as larval forms of an unknown species of *Phyllodromia* or *Ischnoptera*. The species from Abyssinia described but not named by AELUNG (Ann. Mus. Zool. Acad. Imp. St. Petersbourg vol. IX p. 48, fig. 9 (1904)) may be referred to the genus *Apteroblatta* and named after the distinguished Russian entomologist *Adelungi*.

## Fam. **Epilampridæ.**

### Gen. **Calolampra** SAUSS.

#### *Calolampra aptera* SCHULTH.

*Calolampra aptera* de SCHULTHESS SCHINDLER, Ann. Mus. Civ. Gen. (2) Vol. XIX p. 169,  
Pl. II. fig. 2 (1898).

*Kilimandjaro*: Kibonoto. Lower *Meru*: Ngare na nyuki (November—January).

*Usambara*: Tanga. 2 ♂♂, 5 larvae. Previously recorded from Ogaden and Kilimandjaro.

### Gen. **Eustegasta** GERSTAECKER.

A careful examination of several species of this genus has convinced me that the genus is more naturally placed in the family *Epilampridæ* than in the *Perisphæriidæ*. The femora are armed, though sparsely, beneath, the front femora with three to four spines on the anterior margin, the hind femora with two to three on the anterior margin and one to two on the posterior margin, the mid femora are unarmed or else bear one spine on the anterior margin. The supra-anal lamina of the female is produced and generally the apex is cleft, so that it appears sub-bilobate. The facts that the sexes are alike and that the species are viviparous add a little more support to the view, based on structural features, that *Eustegasta* is out of place in the *Perisphæriidæ*.

Pronotum and tegmina marked with rufous . . . . . *pæcila* SCHAUM.

Pronotum and tegmina unicolorous . . . . . *obsoleta* KIRBY.

#### *Eustegasta obsoleta* KIRBY.

. *Eustegasta obsoleta* KIRBY, Ann. Mag. Nat. Hist. (7) vol. V. p. 287, 1900.

*Usambara*: Tanga (June). One female. The species has also been recorded from Nyassaland.

#### *Eustegasta pæcila* SCHAUM.

*Panchlora pæcila* SCHAUM., Ber. Akad. Berlin p. 777 (1853); Peter's Reise Mossamb., Zool. vol. V. p. 109, pl. 7, f. 2 (1862).

*Usambara*: Mombo (June). One female. Previously recorded from Mozambique and Nyassaland.

## Fam. **Blattidæ.**

The old-world genera of Blattidæ with the posterior metatarsus longer than the remaining joints and the tibial spines in three rows may provisionally be distinguished as follows:

1. Tegmina of ♂ reduced, sometimes very short, wings present or absent; of ♀ reduced, sometimes quadrate, never squamiform.

### **Blatta** L.

Type *orientalis* L.

2. Tegmina and wings of ♂ longer than body, no scent-glands opening on 1<sup>st</sup> abdominal tergite, mesonotum and metanotum with short backwardly directed processes. Tegmina of ♀ quadrate, wings absent, penultimate tergite not concealed by antepenultimate tergite.

### **Cartoblatta** gen. n.

Type *pulchra* sp. n.

3. Tegmina and wings of ♂ longer than body, scent-gland opening on 1<sup>st</sup> abdominal tergite, mesonotum and metanotum with long backwardly-directed processes. Tegmina of ♀ squamiform, wings absent, penultimate tergite almost concealed by antepenultimate tergite which is depressed and declivous.

### **Pseudoderopeltis** KRAUSS.

Type *antennata* SAUSS.

4. Tegmina squamiform in both sexes, in ♀ penultimate tergite not concealed by antepenultimate tergite, which is not declivous.

### **Stylopyga** FISCH.

Type *rhombifolia* STOLL.

5. Tegmina and wings longer than the body in both sexes; pronotum trapezoidal, anterior border arcuate, sides deflexed.

### **Periplaneta** BURM.

Type *americana* L.

6. Closely allied to *Periplaneta* but the pronotum elliptical, the anterior border truncate, sides not deflexed.

### **Homalophilpa** STÅL.

Type *ustulata* BURM.

It is not easy to separate females of *Stylopyga* from females of *Pseudoderopeltis*, the declivous 6<sup>th</sup> abdominal tergite almost entirely hiding the 7<sup>th</sup> tergite is perhaps the most distinctive feature of the latter genus, giving the insects a peculiar truncate

appearance when viewed from the side. It is probable that the genus *Blatta* will have to be further subdivided, *orientalis* L., *concinna* HAAN, and *assiniensis* BOL. to take three examples, do not appear to be congeneric.

### Gen. *Paramethana* nov.

Differs from *Dorylaca*<sup>1</sup> STÅL in the size of the second pulvillus of the posterior tarsi which covers the whole second joint, and differs from *Methana* STÅL in the reduction of the tegmina and wings which do not extend beyond the fifth abdominal tergite. Third antennal joint nearly three times as long as the second.

#### *Paramethana robusta* sp. n.

(Pl. 2, fig. 7.)

♀. Dark castaneous, nitid, broadly elliptical. Head castaneous, clypeus and mouth-parts rufo-testaceous, eyes less remote than antennal sockets, antennae fuscous, except the two basal joints which are rufo-testaceous; vertex of head not covered by pronotum. Pronotum broad, trapezoidal, sides deflexed, posterior margin sinuate, castaneous with a central rufo-castaneous mark, or rufo-castaneous with darker castaneous marks. Tegmina short and broad, overlapping considerably, semi-corneous, veins well-marked, mediastinal vein bifurcate, 9 costals, the last three ramose, anal vein deeply impressed, reaching sutural margin at a point one-third from apex. Wings a little shorter than tegmina, flavo-hyaline, apex slightly infuscated. Abdomen piceous above, castaneous below, posterior angles of penultimate tergite strongly produced, its posterior margin slightly produced, sinuate; supra-anal lamina produced, triangular, apex emarginate and deeply cleft. Cerci long, subaeuminate, of thirteen joints. Legs rufo-castaneous, front femora with a complete row of spines on anterior margin beneath; tibial spines in three rows; posterior metatarsus exceeding remaining joints in length, bi-spinulose beneath, pulvilli of remaining joints large occupying the whole extent of the under surface of the joints. Total length 23 mm.; length of tegmina 13 mm.; pronotum 7 mm. × 10,1 mm.

Lower *Meru* (November); 4 ♀♀.

### Gen. *Blatta* L.

#### *Blatta propinqua* sp. n.

Allied to *B. flavilatera* SAUSS. but tegmina in both sexes sub-lobiform.

♂. Dark castaneous. Head with clypeus flavid, ocelli not visible, antennae rufo-castaneous. Pronotum with a broad semicircular band of rufous on each lateral

<sup>1</sup> The type of the genus *Dorylaca* is *brunneri* STÅL, a species closely allied to *flavicincta* HAAN; both species are characterized by peculiar maxillary palpi and by the small size of the second pulvillus of the posterior tarsi; these are good generic characters. In Biol. Centr. Amer. Orth. I. p. 69, 1893, de SAUSSURE and ZEHNTNER re-define the genus *Dorylaca* and transfer to it the apterous species *rhombofolia* STOLL, which is certainly not congeneric with *flavicincta* HAAN and *brunneri* STÅL. This procedure is quite unnecessary. *rhombofolia* is a characteristic species of the genus *Stylopyga*, no useful purpose is served by forcing it into a genus that already includes two species marked by different generic characters.

margin. Tegmina nearly as broad as long, extending to middle of metanotum, their sutural margins failing to meet by half the breadth of the tegmen. Wings absent. Abdomen piceous above, 7<sup>th</sup> tergite slightly produced, its posterior margin sinuate, supra-anal lamina subquadrate, its posterior margin notched; abdomen beneath castaneous at base, piceous at apex, sub-genital lamina broad, extending a little beyond the supra-anal lamina, with two symmetrically disposed styles. Cerci piceous, acuminate. Legs rufous; pulvilli minute, apical.

♀. Resembles ♂ but piceous instead of castaneous. Tegmina relatively shorter and narrower, sub-triangular in shape. Seventh abdominal tergite, more produced; supra-anal lamina produced, cucullate, apex broadly emarginate. Legs darker.

♂. Total length 15 mm.; tegmina 3 mm. × 2,9 mm.; pronotum 5 mm. × 6,1 mm.  
 ♀. " " 18,4 mm.; " 3 mm. × 3 mm.; " 5,1 mm. × 7 "

*Kilimandjaro*: Kibonoto 1,000—1,300 mètres. Lower *Meru*: Masai steppes; 1 ♂, 1 ♀, 4 larvae.

The species closely resembles *Blatta flavilatera* SAUSS. but can at once be distinguished by the very reduced tegmina. The variety *castanea* ADEL. is probably a distinct species, the male has a prominent scent-gland opening on the 1<sup>st</sup> abdominal tergite as in the genus *Pseudoderopeltis* KRAUSS. *B. propinqua* affords a passage from the genus *Blatta* to the genus *Stylopyga* FISCH. *Blatta* at present may be reserved for those species in which the tegmina are not squamiform in both sexes and *Stylopyga* for those species with the tegmina squamiform or absent in both sexes but the discovery of a few more species like *propinqua* would cause this generic distinction to break down.

### Gen. *Stylopyga* FISCH.

#### *Stylopyga hottentota* SAUSS.

*Dorylaea hottentota* DE SAUSSURE, Abhandl. Senckenb. Ges. XXI. p. 578 (1899).

Lower *Meru* (November); 1 ♀.

The species was previously recorded from East Africa e coll. VOELTZKOW. The specimen before me differs slightly from DE SAUSSURE's description but I believe that it must be referred to that species. There are eight species of *Stylopyga* occurring in Africa which can only be distinguished from one another with great difficulty and I believe that some of the species can be sunk as synonymous with others. The species are:

- |   |                       |
|---|-----------------------|
| 1. <i>S. aethiopica</i> SAUSS.                                | Gaboon.               |
| 2. <i>S. manca</i> GERST.                                     | Cameroons.            |
| 3. <i>S. anthracina</i> GERST.                                | Cameroons.            |
| 4. <i>S. spinulifera</i> KRAUSS.                              | San Thomé, W. Africa. |
| 5. <i>S. hottentota</i> SAUSS.                                | E. Africa.            |
| 6. <i>S. brancsiki</i> n. n. (= <i>S. anthracina</i> BRANCS.) | Zambesi.              |
| 7. <i>S. senecta</i> REHN.                                    | Zululand.             |
| 8. <i>S. tetra</i> WLK.                                       | Natal.                |

*Stylopyga spinulifera* KR. and *senecta* REHN can be distinguished by the form of the supra-anal lamina from all the other species; it is probable however that *senecta* REHN is synonymous with *tetra* WLK.; *anthracina* GERST. is possibly the male of *manca* GERST. In *aethiopica* SAUSS. the posterior angles of the last four segments are produced to form backwardly projecting teeth and the two last of these are reflected upwards; in *manca* GERST. the posterior margins of the abdominal tergites are furnished with a row of fine tubercles or plications, but both these characters may be present in both species. ADELUNG has recorded *manca* GERST. from Abyssinia with some doubt, the occurrence of West African species of insects in East Africa is not unknown and I should not be surprised to learn that the number of African species of *Stylopyga* could be reduced to two or three widely distributed forms. *S. rhombifolia* STOLL. has been recorded from Wanga, East Africa (VON DER DECKEN).

### Gen. *Cartoblatta* nov.

♂ allied to *Periplaneta* BURM. but the pronotum transversely elliptical, anterior border truncate, posterior border slightly produced.

Tegmina and wings considerably longer than the body. Mesonotum and metanotum without long backwardly-directed processes. No scent-gland opening on first abdominal tergite.

♀ with the tegmina short, quadrate, not covering the first abdominal tergite. Sixth and seventh abdominal tergites slightly declivous, seventh tergite not covered by the sixth.

### *Cartoblatta pulchra* sp. n.

(Pl. 2. fig. 4.)

♂. Head pale testaceous, the vertex, a curved band between the eyes, a band at base of clypeus castaneous, antennae fuscous. Pronotum pale testaceous, a complex lyrate mark on the disc and a few points on the lateral margins piecous; disc with an anterior and two lateral impressions. Tegmina, anterior part of the wings and the veins castaneous. Abdomen above flavo-testaceous with fusco-castaneous markings laterally, beneath testaceous heavily marbled with fusco-castaneous; supra-anal lamina quadrate, posterior angles rounded; sub-genital plate produced beyond the supra-anal lamina, posterior angles rounded, styles long, curved, arising from notches in the sides of the sub-genital plate. Cerci moderate. Legs testaceous, blotched with fusco-castaneous; posterior metatarsus equal in length to the remaining joints. 3<sup>rd</sup> joint not spined, pulvilli apical but rather large.

♀. Similar to ♂ but head less pale testaceous, the vertex not entirely castaneous, the curved band between the eyes narrower. Pronotum trapezoidal, posteriorly obtusely angled, anteriorly truncate, laterally deflexed, flavo-testaceous with more numerous castaneous points. Scutellum exposed. Tegmina castaneous, quadrate, posterior margin slightly concave. Abdomen flavo-testaceous heavily marbled with fusco-castaneous, posterior margin of 7<sup>th</sup> tergite sinuate, supra-anal lamina narrow,

produced, its apex cleft, genital valves castaneous. Cerci rufous, directed upwards. Posterior metatarsus rather shorter than remaining joints, pulvilli large.

♂. Total length 29 mm.; length of body 20 mm.; length of tegmina 25 mm.; pronotum 4,6 mm. × 7 mm.

♀. Total length 21 mm.; length of tegmina 5,4 mm.; pronotum 5 mm. × 8,2 mm.

*Kilimandjaro*: Kibonoto 1,300—1,900 mètres (March to May); 2 ♂♂, 1 ♀, 8 ♀ larvae.

It is probable that *Stylopyga hova* SAUSS. from Madagascar also belongs to this genus.

### Gen. *Pseudoderopeltis* KRAUSS.

#### *Pseudoderopeltis fulvornata* sp. n.

(Pl. 2, fig. 9.)

♂. Head fusco-castaneous, ocelli, genae and mouth-parts testaceous, apical joint of maxillary palpi infuscated, basal two joints of antennae testaceous, remainder fuscous. Pronotum with two oblique impressions, castaneous, lateral margins testaceous-hyaline, disc with an irregular flavo-testaceous mark on either side of the middle line. Meso- and metanotum with backwardly directed processes, the latter long and slender. Tegmina rufo-castaneous, extending considerably beyond the end of the abdomen. Abdomen above testaceous at base, castaneous at apex, supra-anal lamina quadrate, its posterior margin slightly concave, testaceous with a central castaneous macula. Abdomen beneath castaneous, the disc rufo-castaneous. Cerci moderate. Legs castaneous, posterior metatarsus very long, pulvilli minute, second joint spined beneath.

Total length 23,5 mm.; length of body 16,9 mm.; length of tegmina 20 mm.; pronotum 4 mm. × 5,9 mm.

*Kilimandjaro*: Lower Kibonoto (February), 2 ♂♂.

Apparently allied to *Periplaneta brunneriana* SCHULTH.<sup>1</sup> from Somaliland, a true *Pseudoderopeltis*, and to *P. gildessa*<sup>2</sup> ADEL. from Gallaland, but much smaller than either.

#### *Pseudoderopeltis petrophila* sp. n.

(Pl. 2, fig. 5—6.)

Allied to *P. saussurei* ADEL. ♂. Differs from *P. saussurei* ADEL.<sup>3</sup> in the following points: Head entirely black, nitid, except for the testaceous ocelli and rufous clypeus. Pronotum with the lateral yellow fasciae rather broader and extending to the posterior margin. Tegmina and anterior part of the wings dark castaneous. Abdomen above piceous, castaneous at base, beneath, entirely piceous, nitid.

♀. Nitid, piceous ornamented with yellow; allied to *P. spectabilis* ADEL.<sup>4</sup> (= *P. saussurei* ♀) but head entirely black, except for ocelli and clypeus, which are as

<sup>1</sup> Ann. Mus. Civ. Gen. XXXIX p. 167 pl. 2, f. 1 (1898).

<sup>2</sup> Ann. Mus. Zool. St. Petersburg VIII, p. 314 (1903).

<sup>3</sup> Ann. Mus. Zool. de l'Acad. Imp. Sci. St. Petersburg VIII, p. 316 (1903).

<sup>4</sup> l. c. IX, p. 467 (1905).

in the male; vertex rather flattened, rugose, face with two depressions between the antennal sockets, lower face transversely wrinkled. Pronotum with lateral vittae and two marks on the posterior fifth of the disc yellow, these marks are in some specimens joined to the lateral vittae. Tegmina piceous, lobiform, only just extending beyond the mesonotum, their apices rounded. Mesonotum with two irregular yellow marks on the disc. Metanotum with an irregular transverse yellow band. The five basal abdominal tergites each with a broad yellow band occupying the greater part of their surfaces; 6<sup>th</sup> tergite enlarged, concavely depressed yellow; 7<sup>th</sup> tergite short, triangularly produced, yellow; supra-anal lamina tectiform, carinate, apex emarginate, yellow with a black line on the margin. Cerei and abdomen beneath piceous.

Measurements of types:

♀. Total length 31 mm.; length of body 23 mm.; length of tegmina 26 mm.; pronotum 5,6 mm. × 7 mm.

♀. Length of body 22 mm.; length of tegmina 3,5 mm.; pronotum 7 mm. × 10 mm.

A very long series (115 specimens) of both sexes in all stages of growth from *Kibonoto* 1,300—2,000 mètres, Masai steppes and Lower *Meru* in the acacia forest; the youngest larvae were taken in August and September, older larvae in September and October, the adults in November and December. The females and larvae were found in great quantities under stones bestrewing the steppe-country. Two males from Lower *Meru* exhibit slight variations, in one the yellow fasciae of the pronotum are reduced to short and narrow stripes, in another they are absent altogether; the size of the latter specimen is considerably less than that of the type, but I can find no other characters entitling it to separate specific rank and it may safely be regarded as an extreme variation. With this valuable and important series of specimens before me it is now possible for the first time to distinguish the females of the genus *Pseudoderopeltis* from the females of allied genera with certainty; as already stated their characteristic feature is the enlarged sixth abdominal tergite, which is concavely depressed and declivous so that the posterior part of the body in profile view appears as if it was obliquely truncated.

The species described by me from British E. Africa as *Blatta rothschildi*<sup>1</sup> must be referred to *Pseudoderopeltis*, the female exhibits all the characters of the genus and the male is possibly an abnormal specimen, for the tegmina are reduced but the right tegmen is shorter than the left; the other male characters such as the form of the meso- and metanotum and the opening of the scent-gland on the 1<sup>st</sup> abdominal segment conform to the *Pseudoderopeltis* type. *Stylopyga guttata* SAUSS.<sup>2</sup> from Gallaland is possibly also a ♀ *Pseudoderopeltis*.

<sup>1</sup> Ann. Mag. Nat. Hist. (7) XIX. p. 39 (1907).

<sup>2</sup> Ann. Mus. Civ. Gen. XXXV. p. 75 (1895).

Gen. **Periplaneta** BURM.

No examples of this cosmopolitan genus occur in Dr. SJÖSTEDT's collection but two species have been previously recorded from Eastern Africa, viz: *P. americana* L. (VON DER DECKEN) and *P. atricollis* SAUSS. (VOELTZKOW).

Gen. **Deropeltis** BURM.**Deropeltis melanophila** WLK.

*Ischnoptera melanophila* WALKER, Cat. Blatt. Brit. Mus. Suppl., p. 146. 1869.

*Deropeltis madecassa* DE SAUSSURE, Soc. Ent. VI, p. 17 (1891); DE SAUSSURE and ZEHNTNER Grandidier's Hist. de Madagascar, Orth. I, p. 77, pl. 3, ff. 28, 29 (1895).

Lower *Meru*: Meru rain-forest 3,000 m. (Oct.—Dec.). *Usambara*: Tanga; 4 ♂♂, 2 ♀♀ larvae.

Also recorded from Madagascar, Zanzibar and East Africa.

**Deropeltis integerrima** BR.

*Deropeltis integerrima* BRUNNER, Nouv. Syst. des Blatt. p. 245 (1865).

Lower *Meru*: Ngare na nyuki (November—January); 2 ♂♂, 6 ♀♀.

Previously recorded from Zanzibar; there is a specimen in the Paris Museum from Mombasa.

**Deropeltis autraniana** SAUSS.

*Deropeltis autraniana* SAUSSURE, Ann. Mus. Civ. Gen. XXXV, p. 78 (1895).

A long series (96 specimens) from Lower *Meru*, the steppe country and acacia forest (Sept. to Dec.) and *Kilimandjaro*: Kibonoto 1,000—1,300 mètres. Abundant under stones.

The species is very variable in size in both sexes and the smaller species are less nitid than the larger. There are specimens in the Hope Museum, Oxford from Nairobi, British East Africa and the species was recorded by DE SAUSSURE from Gallaland and West Africa; it is very close to *D. wahlbergi* STÅL from S. Africa and may eventually prove to be identical with it. The determination of the species of the genus *Deropeltis* is attended with considerable difficulty, as the number of forms is considerable and a good proportion have been described from one sex alone. The following tables will perhaps render more easy the tedious task of hunting through literature, whenever one of the more obscure species of the genus has to be identified.

**Males.**

- (32) 1. Fuscous or piceous species.
- (13) 2. Head and legs or legs alone different in colour to rest of body.
- (8) 3. Head and legs rufous or castaneous.



- (5) 4. Outer margin of tegmina sinuate . . . . . *erythrocephala* FAB. (S. Africa).  
 (4) 5. Outer margin of tegmina not sinuate.  
 (7) 6. Large species . . . . . *dmitriewi* ADEL. (Abyssinia).  
 (6) 7. Small species . . . . . *gracilis* BURM. (S. Africa).  
 (3) 8. Legs only different in colour to rest of body.  
 (10) 9. Large species . . . . . *erythropeza* ADEL. (Abyssinia).  
 (9) 10. Small species.  
 (12) 11. Legs red . . . . . *intermedia* BR. (Natal).  
 (11) 12. Legs testaceous . . . . . *sculpturata* KRAUSS. (Guinea).  
 (2) 13. Head and legs concolorous with rest of body.  
 (15) 14. Densely pubescent . . . . . *integerrima* BR. (Zanzibar, E. Africa).  
 (14) 15. Not densely pubescent.  
 (21) 16. Small species, body-length not exceeding 20—22 mm.  
 (20) 17. Tegmina considerably exceeding the body.  
 (19) 18. Pronotum opaque, finely punctate . . . . . *wahlbergi* STÅL. [= *atra* BR.] (S. Africa).  
 (18) 19. Pronotum nitid . . . . . *autraniana* SAUSS. (Gallaland, E. and W. Africa).  
 (17) 20. Tegmina scarcely exceeding the body . . . . . *kuchovskii* ADEL. (Abyssinia).  
 (16) 21. Larger species.  
 (29) 22. Tegmina considerably exceeding the body.  
 (24) 23. Pronotum relatively very small . . . . . *adelungi* nom. n. [= *gracilis*, ADEL.] (Abyssinia).  
 (23) 24. Pronotum relatively larger.  
 (26) 25. Posterior margin of mesonotum with long triangular processes . . . . . *pallidipennis* ADEL. (Abyssinia).  
 (25) 26. Posterior margin of mesonotum without these processes.  
 (28) 27. Body-length 22 mm. . . . . *nigrita* SAUSS. (Somaliland).  
 (27) 28. Body-length 29 mm. . . . . *melanophila* WLK. (Madagascar, E. Africa).  
 (22) 29. Tegmina not exceeding body by much.  
 (31) 30. Body-length 23 mm. . . . . *barbeyana* SAUSS. (Somaliland).  
 (30) 31. Body-length 36,5 mm. . . . . *tullbergi* BORG (Cameroons).  
 (1) 32. Ferruginous or testaceous species.  
 (34) 33. Pronotum piceous . . . . . *bucana* KARSCH. (Cameroons).  
 (33) 34. Pronotum not piceous.  
 (36) 35. Pronotum with three impressions on disc . . . . . *triimpressa* KRAUSS. (Guinea).  
 (35) 36. Pronotum not as above . . . . . *negus* ADEL. (Abyssinia).

## Females.

- (24) 1. Fuscous or piceous species.
- (23) 2. Without rufous fasciae on the pronotum.
- (8) 3. Head and legs castaneous or rufous.
- (5) 4. Castaneous . . . . . *erythropeza* ADEL.
- (4) 5. Rufous.
- (7) 6. 6<sup>th</sup> and 7<sup>th</sup> abdominal tergites with a fulvous macula on either side . . . . . *dichroa* GERST. (Gold Coast).
- (6) 7. Abdominal tergites uniformly coloured . . . *erythrocephala* FAB.
- (3) 8. Head and legs concolorous with rest of body.
- (10) 9. Densely pubescent . . . . . *integerrima* BR.
- (9) 10. Not densely pubescent.
- (20) 11. Angles of thoracic tergites strongly produced backwards.
- (19) 12. Species exceeding 24 mm. in length.
- (18) 13. Posterior margin of pronotum straight.
- (17) 14. Posterior angles of abdominal tergites spiniform.
- (16) 15. 38 mm. in length . . . . . *robusta* GERST. (Cameroons).
- (15) 16. 25 mm. in length . . . . . *gaboonica* REHN (Gaboon).
- (14) 17. Posterior angles of abdominal tergites not spiniform . . . . . *melanophila* WLK. [= *speiseri* BRANCS.]
- (13) 18. Posterior margin of pronotum sinuate. . . . *schweinfurthi* SAUSS. (Somaliland).
- (12) 19. Smaller species . . . . . *nigrita* SAUSS.
- (11) 20. Angles of thoracic segments not strongly produced backwards. . . . . *kachovskii* ADEL.
- (22) 21. Posterior margin of pronotum sinuate . . . *barbeyana* SAUSS.
- (21) 22. Posterior margin of pronotum straight . . . *carbonaria* GERST. (W. Africa).  
*autraniana* SAUSS.
- (2) 23. With rufous fasciae on the margins of the pronotum . . . . . *paulinoi* BOL. (Angola).
- (1) 24. Ferruginous species . . . . . *trimpresa* KRAUSS.

As our knowledge of the genus extends, it will probably be found that some of these names are synonymous, e. g. *D. robusta* GERST. may prove to be the female of *D. tullbergi* BORG. DE SAUSSURE gives the range of *D. autraniana* as West Africa (Abetifi), Somaliland and Gallaland; from the descriptions alone I find it impossible to distinguish de SAUSSURE'S species from *D. carbonaria* GERST. The following species which have been recorded by various authors as of the genus *Deropeltis*, either belong to other genera or else are of uncertain position:

<i>Blatta capensis</i> THUNB.	= <i>Deropeltis erythrocephala</i> FAB.
<i>Euryzosteria delalandii</i> SAUSS.	= <i>Deropeltis erythrocephala</i> FAB.
<i>Perisphaeria verticalis</i> BURM.	= ? <i>Deropeltis</i> .
<i>Periplaneta orba</i> STÅL.	= <i>Pseudoderopeltis</i> .
<i>Periplaneta albilatera</i> STÅL.	= <i>Pseudoderopeltis</i> .
<i>Periplaneta caffra</i> STÅL.	= <i>Periplaneta caffra</i> .
<i>Kakerlac brevicollis</i> SERV.	= ? <i>Pseudoderopeltis</i> .
<i>Ischnoptera longipennis</i> WLK.	} = <i>Pseudoderopeltis</i>
<i>Ischnoptera juncea</i> SAUSS.	
<i>Ischnoptera similis</i> SAUSS.	
<i>Deropeltis longipennis</i> SAUSS.	
<i>Deropeltis antennata</i> SAUSS.	
<i>Polyzosteria capensis</i> SAUSS.	} = <i>Pseudoderopeltis</i> .
[= <i>Deropeltis flavomarginata</i> BR.]	
<i>Polyzosteria meridionalis</i> SAUSS.	} = <i>Blatta</i> .
[= <i>Deropeltis bivittata</i> BR.]	
[= <i>Deropeltis distanti</i> KIRBY]	
<i>Ischnoptera macra</i> STÅL.	= <i>Perisphaeriinae</i> .
<i>Deropeltis burmeisteri</i> SAUSS.	} ? <i>nomina nuda</i> , only mentioned in a synoptical key
<i>Deropeltis peringueyi</i> SAUSS.	

## Fam. Panchloridæ.

### Gen. Leucophaea BR.

#### *Leucophaea* sp.

*Kilimandjaro*: Kibonoto 1,300—1,900 (November). 9 larvae (♂ and ♀).

These are not the larvae of *L. surinamensis* L. being considerably larger; it is in the highest degree probable that they belong to an undescribed species. *Surinamensis* has been recorded from Wanga (VON DER DECKEN).

### Gen. Panchlora BURM.

#### *Panchlora camerunensis* BORG.

*Panchlora camerunensis* BORG. Bih. K. Svenska Vet. Akad. Afd. IV, No. 10, p. 24, 1902.

*Kilimandjaro*: Kibonoto 1,000—1,900 mètres (November, March), 8 ♂♂, 8 ♀♀, 1 larva.

This appears to be the same species as that described by BORG from the Cameroons, also collected by Dr. SJÖSTEDT; the East African specimens are a trifle larger but otherwise appear to be identical.

Gen. **Nauphoeta** BURM.**Nauphoeta cinerea** OLIV.

*Blatta cinerea* OLIVIER, Enc. Méth. Ins. IV. p. 314, n. 3, 1789.

*Kilimandjaro* (Sept.); Kibonoto 1,000—1,900 mètres (May); *Usambara*: Mombo (June): 2 ♂♂, 4 ♀♀, 3 larvae.

A widely distributed species.

Gen. **Gyna** BR.**Gyna vetula** BR.

*Gyna vetula* BRUNNER V. WATTENWYL, Nouv. Syst. des Blatt. p. 267 (1865).

In the Mkulumusi-caves at *Tanga* (July), 1 ♂, 1 ♀, 17 larvae.

Previously recorded from Mombasa (VON DER DECKEN) and the East coast of Africa.

It seems quite certain that the genera *Trichomeva* KIRBY<sup>1</sup> and *Apotrogia* KIRBY<sup>2</sup> are founded on immature specimens of species of *Gyna*. The larvae collected by Dr. SJÖSTEDT are certainly most closely allied to *Trichomeva insignata* KIRBY; they are preserved in alcohol together with one adult male in bad condition and were probably taken altogether from under one shelter. The backwardly produced angles of the meso- and metanotum proclaim the immature condition of these specimens as well as of *T. insignata* KIRBY and *Apotrogia angolensis* KIRBY. I find moreover that the structure of the maxillary and labial palpi, the form of the coxae and the arrangement of speines on the tibiae is identical in the larvae and adults. The palpi in the genus *Gyna* are highly characteristic, being very slender and rather long, and the coxae are provided with a peculiar curved flange, situated on the outer posterior angle; the front tibiae in the larvae are shorter than they are in the adults, but this feature is probably associated with fossorial habits, abandoned when the insects become winged adults. The arolium between the tarsal claws does not develop until the larvae are nearly full-grown and it is then smaller than in the adult. The pronotum is far less backwardly produced in the larvae than in the adults, as is also the case with larval Epilamprides, larvae of *Panchlora* and of *Rhyparobia*. Finally the larvae are much more heavily marked with castaneous than are the adults and there are rows of minute tubercles on the dorsal tergites.

Gen. **Phenacisma** KARSCH.

The type of the genus, *P. peltata* KARSCH has been recorded from Mombasa.

<sup>1</sup> Ann. Mag. Nat. Hist. (6) XVIII. p. 257 (1896).

<sup>2</sup> ibidem (7) V. p. 281 (1900).

Fam. **Corydiidae.**Gen. **Anacompsa** nov.

Finely pubescent. Eyes approximate. Antennae filiform, shorter than body. Pronotum cucullate, covering vertex of head, small, sides deflexed, posterior border arcuate. Tegmina and wings elongate, extending considerably beyond the apex of the abdomen; costal veins few and irregular, discoidal area of tegmina reticulated, axillary veins obsolescent; posterior part of wing ample, median vein simple, ulnar vein ramose at extremity. Supra-anal lamina of ♂ transverse. Cerci moderate. Tibiae weakly armed, spines on posterior pair triseriate with 4 apical calcaria, 4 apical calcaria on front pair. Tarsal claws with arolia.

In general appearance the genus resembles *Latindia* STÅL but is distinguished from that and its allies, *Hemilatindia* SAUSS. and *Paralatindia* SAUSS. by the triseriate arrangement of the tibial spines. From *Holocompsa* BURM. and *Hypercompsa* SAUSS. it is distinguished by the long tegmina of uniform texture; *Ipisoma* BOL. is characterised *inter alia* by the reduced tegmina. It is probable that a considerable number of species of these fragile little cockroaches are still unknown to science, they are rare in collections but probably because they have been neglected by most collectors in favour of larger and more conspicuous forms.

**Anacompsa cucullata** sp. n.

(Pl. 2, fig. 10 and 11, Pl. 3, fig. 9.)

♂. Head dark castaneous, mouth parts testaceous; frons not swollen, ocelli large. Antennae fuscous, filiform, of 30 joints, joints longer than broad and increasing in length distally. Pronotum rufo-testaceous, disc with two oblique impressions. Tegmina long, overlapping strongly, finely pubescent, membranous, rufo-testaceous with a broad humeral vitta extending to half the total length, fuscous; mediastinal vein with a few incomplete branches, 6 costal veins, their ends not reaching the anterior margin of the tegmen, anterior ulnar ramose at its apex, posterior ulnar ramose, transverse anastomoses uniting the branches, anal vein curved reaching the sutural margin at a point one-fifth of the total length, axillary veins obsolescent. Wings hyaline, flavo-testaceous on the sutural margin, mediastinal vein simple, 6 irregular costals, ulnar vein ramose at its extremity, posterior field ample larger than the anterior field. Abdomen castaneous, flavo-testaceous at base, supra-anal lamina transverse, sub-genital small, right style absent, left style represented by a flattened lobe, the origin of which is hidden by the penultimate sternite. Cerci moderate, testaceous, segmentation obscure. Legs testaceous; front tibiae with only 4 spines on the anterior border, 2 being apical, 3 on the posterior border, 2 being apical; spines on mid- and hind-tibiae triseriate; front femora with no genicular spine, formula of apical spines  $\frac{1}{0}$ ,  $\frac{1}{0}$ ,  $\frac{1}{0}$ .

Total length 10,5 mm.: length of body 7,5 mm.; length of tegmina 9 mm.; pronotum 2 mm.  $\times$  2,9 mm.

Lower *Meru*: Ngare na nyuki (January); 2 ♂♂.

### Gen. *Sphecophila* SHELF.

#### *Sphecophila termitum* sp. n.

(Pl. 3, figs. 1—2.)

♂. Fulvous, rufo-fimbriate. Head with vertex not covered by pronotum, antennae consisting of 30 joints, similar in structure to those of *S. polybiarum* mihi,<sup>1</sup> frons swollen, ocelli minute, eyes much reduced. Body above with a minute recumbent pubescence, margins rufo-fimbriate. Posterior margins of pronotum and of mesonotum slightly obtusely angled, postero-lateral angles of meso- and meta-notum backwardly produced. Ten abdominal tergites visible, 8<sup>th</sup> and 9<sup>th</sup> very narrow almost concealed beneath the 7<sup>th</sup>, supra-anal lamina quadrangularly produced, angles rounded, posterior margin slightly excised; cerci short, one-jointed, acuminate; seven abdominal sternites visible, subgenital lamina sub-triangularly produced, rounded, with a pair of slender styles, podical plates prominent, tumid. Legs as in *S. polybiarum*, except that there is an anterior apical spine on the front femora, genicular spines of first and second femora very stout and long.

Total length 7,1 mm.; pronotum 2,1 mm.  $\times$  4,1 mm.

*Kilimandjaro*: Kibonoto, March 1906. 2 ♂♂ from nest of *Termes bellicosus*.

It is certainly surprising to find a species of cockroach symbiotic with termites in E. Africa, congeneric with a South American species living in a wasp's nest; I have utterly failed to find, however, any characters entitling the former species to separate generic rank from the latter. The large size and the brilliant fulvous colour of the African species readily serve to distinguish it from *S. polybiarum*, but in both species the structure of the head, eyes, mouth-parts, form of the body, armature of the legs, shape of the terminal abdominal scutes is closely similar. Dr. SJÖSTEDT's specimens have been preserved in alcohol and the abdominal segments have become somewhat distended, so that the podical plates appear very prominently, but they are also clearly visible in dried specimens of *polybiarum* and their prominence may possibly be regarded as a character of the genus. Further collections of cockroaches symbiotic with other insects will reveal perhaps some day the wide distribution of this genus. At present the species described here is the only cockroach living in company with social insects that has yet been recorded from the Old World.

### Gen. *Euthyrrhapha* BURM.

The cosmopolitan species *E. pacifica* Coq. has been recorded from Wanga (VON DER DECKEN).

<sup>1</sup> Trans. Ent. Soc. London 1906, p. 518.

Fam. **Oxyhaloidæ.**Gen. **Oxyhaloa** BR.

Key to E. African species.

1. Tegmina smooth . . . . . *O. ferreti* REICHE and FAIRM.
- 1'. Tegmina with minute recumbent pubescence.
  2. Tegmina attaining end of abdomen . . . . . *O. deusta* THUNB.
  - 2' Tegmina not attaining end of abdomen. . . . . *O. variabilis* SHELF.

**Oxyhaloa variabilis** SHELF.

*Oxyhaloa variabilis* SHELFORD, Ann. Mag. Nat. Hist. (7) vol. 19, p. 41, 1907.

*Kilimandjaro*: Kibonoto 1,000—1,300 mètres (Feb., Apr., Nov.); 3 ♀. Previously recorded from the interior of Djibouti. The species is very close to *O. deusta* THUNB. (= *fulviceps* BURM.), which together with *O. ferreti* REICHE and FAIRM. has been recorded from Lake Jipe at Kilimandjaro (VON DER DECKEN).

Fam. **Perisphaeriidæ.**Gen. **Gynopeltis** GERST.**Gynopeltis cryptospila** WLK.

*Polyphaga cryptospila* WALKER, Cat. Blatt. Brit. Mus., p. 15 (1868).

*Gynopeltis picta* GERSTAECKER, Arch. Naturg. XXXV, p. 208 (1869); Von der Decken's Reisen in Ost-Afrika, III (2). p. 9, pl. 1, ff. 1, 2 (1873).

*Kilimandjaro* »Mischwald»; Kibonoto 1,000—1,900 mètres; Steppe country. *Meru* 3,000 mètres. 4 ♂♂, 10 ♀♀. Previously recorded from Endara and Mosambique.

Gen. **Derocalymma** BURM.

The two East African species can be distinguished as follows:

**Males.**

1. Body-length 19 mm., antennae entirely fuscous . . . . . *porcellio* GERST.
- 1'. Body length 15 mm., antennae with broad pale annulus . . . *lampyrina* GERST.

**Females.**

1. Length of pronotum : breadth of pronotum : : 550 : 935; antennae entirely fuscous . . . . . *porcellio* GERST.
- 1'. Length of pronotum : breadth of pronotum : : 550 : 1050; antennae with broad pale annulus at base and at apex . . . . . *lampyrina* GERST.

***Derocalymma porcellio* GERST.**

*Derocalymma porcellio* GERSTAECKER Arch. Naturg. XXXV, p. 207 (1869); Von der Decken's Reisen in Ost-Afrika, III (2), p. 7, pl. 1, f. 3 (1873).

Lower *Meru*: Ngare na nyuki. *Kilimandjaro*: Kibonoto 1,000—1,300 mètres (Sept. to Jan.); 19 ♀♀ adult and larval. Previously recorded from Lake Jipe at Kili-*mandjare* and *Uru*, East Africa.

***Derocalymma lampyrina* GERST.**

*Derocalymma lampyrina* GERSTAECKER, ll. cc., p. 207 (1869); p. 8 (1873).

*Usambara*: Mombo (June); 3 ♀♀. Previously recorded from between Lake Jipe and the Bura Mts.

**Gen. *Cyrtotria* STÅL [= *Stenopilema* SAUSS.]**

Dr. SJÖSTEDT has kindly sent to me for examination a number of STÅL's types of Blattidæ. Amongst these are the male and female types of *Cyrtotria gibbicollis* STÅL and they prove to be congeneric with the species included by de SAUSSURE and ZEHNTNER (Rev. Suisse Zool. III, p. 25 (1895) in the genus *Stenopilema*. In the genus *Cyrtotria* the distinguished Swiss orthopterists place *Derocalymma dispar* BURM. *Cyrtotria macra* STÅL and with some doubt *Perisphaeria affinis* BURM.; the first of these is certainly not congeneric with *C. gibbicollis* STÅL and a new genus must be created for its reception; the type is in BRUNNER's collection and has been well described so that it is easily recognisable, the other two species are of uncertain systematic position. *Stenopilema* SAUSS. of course sinks as a synonym of *Cyrtotria* STÅL. It may be noted here that *Perisphaeria linearis* WALK. is synonymous with *Cyrtotria gibbicollis* STÅL.

***Cyrtotria capucina* GERST.**

*Derocalymma capucina* GERSTAECKER. Arch. Naturg. XXXV, p. 207 (1869); Von der Decken's Reisen in Ost-Afrika, III (2), p. 8, pl. 1, f. 4 (1873).

♂. Piecous. Eyes contiguous; antennae fuscous, basal joints paler; ocelli and mouth parts testaceous. Pronotum with margins castaneous, disc strongly punctate with a few smooth interspaces, anteriorly earinate. Tegmina dark castaneous in basal fourth, remainder testaceous, marginal area at base testaceous, eight longitudinal discoidal sectors. Wings hyaline, veins testaceous, ulnar vein sending one branch to apex of wing, six incomplete branches to dividing vein. Supra-anal lamina quadrate, its posterior margin slightly concave; sub-genital lamina irregular, laterally margined with testaceous; cerei and styles testaceous. Coxae castaneous, outwardly margined with testaceous; legs testaceous.

Total length 18,7 mm.; length of body 14 mm.; length of tegmina 15,5 mm.; pronotum 4 mm. × 4 mm.



*Cyrtotria gibbicollis* STÅL is very near this species.

*Kilimandjaro*: Kibonoto 1,000—1,900 mètres, Masai steppes (Aug. to Nov.).  
Lower *Meru*. 2 ♂♂, 33 ♀♀ adult and larval. Previously recorded from Aruseha,  
E. Africa.

*Cyrtotria tuberculata* sp. n.

♀. Allied to *C. capucina* GERST. but the pronotum anteriorly with small scattered tubercles, posteriorly rugose with a few fine punctures. Head more rufous.

Total length 15 mm.; pronotum 5 mm. × 5 mm.

*Kilimandjaro*: Lower Kibonoto (Feb.), 1 ♀.

Gen. *Parasphaeria* BR.

*Parasphaeria* ? *marmorata* sp. n.

(Pl. 2, fig. 8.)

♀. Testaceous, marbled with fuscous. Head only half covered by pronotum; eyes rather small and wide apart, ocelli absent, antennae fuscous, equal to half the body-length; frons rufo-castaneous, face with dark markings. Pronotum trapezoidal, anterior and posterior margins truncate, lateral margins hyaline, disc with a few shallow punctures and with a dark lyrate marking. Tegmina lobiform, testaceous-hyaline; mesonotum and metanotum with scattered punctures and dark markings symmetrically arranged, lateral margins hyaline. Abdominal tergites with scattered shallow punctures laterally bordered with testaceous, their posterior margins broadly olive brown, a central row of fuscous spots and a sub-marginal row on each side of similar markings, in addition some irregular fuscous markings between the central and submarginal rows; 1<sup>st</sup> to 7<sup>th</sup> tergite divided by a transverse suture into a narrow anterior half and a broader posterior half. Supra-anal lamina slightly produced, trigonal; cerci very short, of four joints. Abdominal sternites with submarginal row of castaneous spots, an irregular transverse row of markings on each sternite, except the last which is castaneous on the disc; each sternite except the last is divided like the tergites; sub-genital lamina ample, semiorbicular, extending beyond the supra-anal lamina. Legs short, testaceous, a fuscous line on the outer side of the femora; tibial spines in three rows: posterior metatarsus not equal to the remaining joints; arolia large.

Length 15 mm.; pronotum 3 mm. × 4.5 mm.

Lower *Meru* (Nov.); 1 ♀.

I have been unwilling to place this species in the Neotropical genus *Parasphaeria*, but the unique female specimen certainly presents no features whereby it can be separated therefrom; the exposed head, lobiform tegmina, tibial spines and short tarsi are the distinctive characters of *Parasphaeria* and until the male sex is discovered this species must be allowed a place in that genus.

Table showing the distribution in other parts of Africa of the species of the Kilimandjaro-Meru district.

Kilimandjaro-Meru district	Cosmo- politan	Abyssinia	Somali- land	E. Africa from 5° S. to Zambezi	Africa S. of Zambezi	W. Africa	Mada- gascar
1. <i>Theganopteryx africana</i> . . . . .	.	.	.	✗	.	.	.
2. <i>Theganopteryx saussurei</i> . . . . .	.	.	✗	.	.	✗ <sup>1</sup>	.
3. <i>Mallotoblatta kraussi</i> . . . . .	.	✗	.	.	.	.	.
4. <i>Hololampra aethiopica</i> . . . . .	.	.	.	.	.	.	.
5. <i>Hololampra sjöstedti</i> . . . . .	.	.	.	.	.	.	.
6. <i>Ischnoptera bimaculata</i> . . . . .	.	.	.	✗	.	.	.
7. <i>Ischnoptera incuriosa</i> . . . . .	.	.	.	✗	✗	.	.
8. <i>Ischnoptera neutra</i> . . . . .	.	.	.	✗	.	.	.
9. <i>Phyllodromia germanica</i> . . . . .	✗	.	.	.	.	.	.
10. <i>Phyllodromia bivittata</i> . . . . .	✗	.	.	.	.	.	.
11. <i>Phyllodromia supellectilium</i> . . . . .	✗	.	.	.	.	.	.
12. <i>Phyllodromia zchntneri</i> . . . . .	.	.	.	.	.	.	✗
13. <i>Phyllodromia nigromarginata</i> . . . . .	.	.	.	.	.	.	.
14. <i>Phyllodromia sjöstedti</i> . . . . .	.	.	.	.	.	.	.
15. <i>Phyllodromia insignis</i> . . . . .	.	.	.	.	.	.	.
16. <i>Phyllodromia testacea</i> . . . . .	.	.	.	.	.	.	.
17. <i>Phyllodromia trigonalis</i> . . . . .	.	.	.	✗	.	.	.
18. <i>Ceratinoptera bimaculata</i> . . . . .	.	.	.	.	.	.	.
19. <i>Ceratinoptera castanea</i> . . . . .	.	.	.	.	.	.	.
20. <i>Ceratinoptera sjöstedti</i> . . . . .	.	.	.	.	.	.	.
21. <i>Ceratinoptera variabilis</i> . . . . .	.	.	.	.	.	.	.
22. <i>Ceratinoptera perpulchra</i> . . . . .	.	.	.	.	.	.	.
23. <i>Ceratinoptera ferruginea</i> . . . . .	.	.	.	✗	.	.	.
24. <i>Ceratinoptera ovata</i> . . . . .	.	.	.	.	.	.	.
25. <i>Ceratinoptera variegata</i> . . . . .	.	.	.	+	.	.	.
26. <i>Ceratinoptera dimidiata</i> . . . . .	.	.	.	.	.	.	.
27. <i>Temnopteryx abyssinica</i> . . . . .	.	✗	.	✗	.	.	.
28. <i>Temnopteryx ectobioides</i> . . . . .	.	.	.	.	.	.	.
29. <i>Temnopteryx caffra</i> . . . . .	.	.	.	✗	.	.	.
30. <i>Temnopteryx affinis</i> . . . . .	.	.	.	.	.	.	.
31. <i>Temnopteryx rufa</i> . . . . .	.	.	.	.	.	.	.
32. <i>Loboptera nitida</i> . . . . .	.	.	.	.	.	.	.
33. <i>Apteroblatta perplexa</i> . . . . .	.	.	.	.	.	.	.
34. <i>Calolampra aptera</i> . . . . .	.	.	✗	.	.	.	.
35. <i>Eustegasta obsoleta</i> . . . . .	.	.	.	✗	.	.	.
36. <i>Eustegasta pacila</i> . . . . .	.	.	.	✗	.	.	.
37. <i>Paramethana robusta</i> . . . . .	.	.	.	.	.	.	.
38. <i>Blatta propinqua</i> . . . . .	.	×	×	.	.	.	.
39. <i>Stylopyga hottentota</i> . . . . .	.	.	.	✗	.	.	.
40. <i>Stylopyga rhombifolia</i> . . . . .	✗	.	.	.	.	.	.
41. <i>Cartoblatta pulchra</i> . . . . .	.	.	.	.	.	.	.

<sup>1</sup> An cross (×) denotes a closely allied species or topomorph.

Kilimandjaro-Mern District	Cosmo- politan	Abyssinia	Somali- land	E. Africa from 5° S to Zambezi	Africa S. of Zambezi	W. Africa	Madag- ascar
42. <i>Pseudoderopeltis fulvornata</i> . . . . .	.	.	.	.	.	.	.
43. <i>Pseudoderopeltis petrophila</i> . . . . .	.	.	.	.	.	.	.
44. <i>Periplaneta americana</i> . . . . .	×	.	.	.	.	.	.
45. <i>Periplaneta atricollis</i> . . . . .	.	.	.	+	.	.	.
46. <i>Deropeltis melanophila</i> . . . . .	.	.	.	×	.	.	×
47. <i>Deropeltis integerrima</i> . . . . .	.	.	.	+	.	.	.
48. <i>Deropeltis australiana</i> . . . . .	.	×	.	×	.	×	.
49. <i>Panchlora camerunensis</i> . . . . .	.	.	.	.	.	×	.
50. <i>Nauphaeta cinerea</i> . . . . .	×	.	.	.	.	.	.
51. <i>Gyna vetula</i> . . . . .	.	.	.	×	.	.	.
52. <i>Phenacisma peltata</i> . . . . .	.	.	.	.	.	.	.
53. <i>Anacompsa cucullata</i> . . . . .	.	.	.	.	.	.	.
54. <i>Sphecochyla termitium</i> . . . . .	.	.	.	.	.	.	.
55. <i>Euthyrrhapha pacifica</i> . . . . .	+	.	.	.	.	.	.
56. <i>Oxyhaloa deusta</i> . . . . .	.	.	.	×	×	.	.
57. <i>Oxyhaloa ferreti</i> . . . . .	.	×	.	.	.	.	.
58. <i>Oxyhaloa variabilis</i> . . . . .	.	.	.	+	.	.	.
59. <i>Gynopeltis cryptospila</i> . . . . .	.	.	.	×	×	.	.
60. <i>Derocalymma porcellio</i> . . . . .	.	.	.	×	.	.	.
61. <i>Derocalymma lampyroides</i> . . . . .	.	.	.	.	.	.	.
62. <i>Cyrtotria capucina</i> . . . . .	.	.	.	.	×	.	.
63. <i>Cyrtotria tuberculata</i> . . . . .	.	.	.	.	.	.	.
64. <i>Parasphaeria (?) marmorata</i> . . . . .	.	.	.	.	.	.	.

The above table includes 64 species<sup>1</sup> and a glance at their distribution shows as might have been safely predicted that the closest affinity of this Blattid fauna is found in East Africa between the 5<sup>th</sup> degree of latitude and the Zambezi River; 20 of the species are common to these two areas and I have little doubt that the number would be increased if our knowledge of the cockroaches of the latter area was greater than it is. 7 of the species are cosmopolitan or nearly so and consequently afford no evidence worth taking into consideration. The relationship with the Abyssinian fauna is rather surprisingly remote, but can perhaps be accounted for by the intervention of the xerothermic area of Somaliland and Gallaland between the damp moist regions of Abyssinia and Kilimandjaro; it is significant that there is a much closer relationship between the faunas of these two regions in the case of insects capable of sustained flight such as the Lepidoptera Rhopalocera. Only 4 of the species occur also in Africa south of the Zambesi and the entire absence from the Kilimandjaro region of certain highly characteristic South African genera is in sharp contrast to the fact that all the Abyssinian genera save two<sup>2</sup> are represented in the Kilimandjaro region. The connection with the West African and Mascarene faunas

<sup>1</sup> An immature and undetermined species of *Leucophaea* is omitted.

<sup>2</sup> One of these two is *Paraloboptera*, a genus barely separable from *Loboptera* which is represented by one species in the Kilimandjaro regions.

is of the remotest, but that there should be any species common to regions so widely separated is of much interest. The absence from a given fauna of certain genera and species often affords as much food for speculation and as many points of interest as the presence of others, and this fauna offers no exception to such a statement. The genus *Anaplecta* is moderately represented in West Africa but except for one species from the Soudan is absent in the East. *Epilampra* is well represented in West Africa but is entirely absent from the Eastern and Southern regions, its place being filled to a certain extent by the genus *Gyna*, though this too is poorly represented both in numbers of species and individuals in the Kilimandjaro and Abyssinian regions. *Polyphaga* is a characteristically dry-country genus, its absence from Dr. SJÖSTEDT's collection is therefore not surprising; in Abyssinia, Somaliland and South Africa it is well represented. South Africa may be regarded as the head-quarters of the family *Perisphaeriidae* and such genera as *Aptera*, *Pronaonota*, *Pilema*, *Melanoblatta*, *Hostilia*, *Paciloblatta* and others are peculiar to this region and include some of the most abundant species; in the Kilimandjaro region this sub-family is represented by four genera and six species only, two of the species being represented in Dr. SJÖSTEDT's collection by a good number of individuals, nearly all of the female sex, still it cannot be said that this sub-family is dominant in this region. The most dominant species of the Kilimandjaro fauna, as evidenced by the number of individuals captured, are the species of the genera *Pseudoderopeltis* and *Deropeltis*, and notably *P. petrophila* and *D. autraniana*; these were taken in great abundance at all seasons of the year and at varying altitudes and in a letter Dr. SJÖSTEDT informs me that many were found under the boulders bestrewing the Masai steppe-country.

The discovery of a cockroach symbiotic with termites is of much interest and that this species should be strictly congeneric with a species found in the nest of a social wasp in S. America is a fact that can scarcely be explained on any other supposition than that further collecting will reveal the wide spread distribution of the genus throughout the tropics.

PLATE 2.

Plate 2.

- Fig. 1. *Holotampra ethiopica* SHELF. n. sp. ♀.  
 » 2. *Ceratinoptera perpulchra* SHELF. n. sp. ♀.  
 » 3. *Apteroblatta perplexa* SHELF. n. sp. ♀.  
 » 4. *Cartoblatta pulchra* SHELF. n. sp. ♀.  
 5. *Pseudoderopeltis petrophila* SHELF. n. sp. ♀.  
 6. » » » ♀ from the side.  
 7. *Paramethana robusta* SHELF. n. sp. ♀.  
 » 8. *Paraspheria marmorata* SHELF. n. sp. ♀.  
 9. *Pseudoderopeltis fulvornata* SHELF. n. sp. ♂.  
 10. *Anacompsa cucullata* SHELF. n. sp., wing.  
 » 11. » » » tegmen.  
 » 12. *Temnopteryx ectobioides* SHELF. n. sp. ♂; apex of abdomen, ventral view.  
 » 13. *Apteroblatta perplexa* SHELF. n. sp. ♂; » » » » »  
 » 14. *Phyllodromia testacea* SHELF. n. sp. ♂; » » » » dorsal »




PLATE 3.



## Plate 3.

- Fig. 1. *Sphcophila termitium* SHELF n. sp. ♂; Lg 7,1 mm.  
 » 2. » » » ♂; side view of head.  
 » 3. *Mallotoblatta kraussi* ADEL. ♂; apex of abdomen, dorsal view.  
 » 4. *Hololampra aethiopica* SHELF. n. sp. ♂; apex of abdomen, dorsal view.  
 » 5. » » » ♂; titillator of penis.  
 » 6. *Ischnoptera incuriosa* SAUSS. ♂; apex of abdomen, dorsal view.  
 » 7. » » » ♂; » » » , ventral view.  
 » 8. *Phyllodromia insignis* SHELF. n. sp. ♂; apex of abdomen, ventral view.  
 » 9. *Anacompsa cucullata* SHELF. n. sp. ♂; apex of abdomen, ventral view.  
 » 10. *Ischnoptera bimaculata* GERST. ♂; apex of abdomen; dorsal view.  
 » 11. *Phyllodromia supellectilium* SERV. ♂; apex of abdomen, dorsal view.  
 » 12. » *nigromarginata* SHELF. n. sp. ♂; apex of abdomen, dorsal view.  
 » 13. *Ceratinoptera perpulchra* SHELF. n. sp. ♂; apex of abdomen, dorsal view.  
 » 14. *Hololampra aethiopica*. ♂; wing.  
 » 15. *Ischnoptera bimaculata* GERST. ♂; apex of abdomen, ventral view.  
 » 16. *Ceratinoptera sjöstedti* SHELF. n. sp. ♂; apex of abdomen, ventral view.  
 » 17. » » » ♂; titillator of penis.

